

Running Title: RIRS Safe and Efficient in Previous Kidney Stone Surgery

Is RIRS Safe and Efficient In Patients With Kidney Stones Who Had Previous Open ,Endoscopic,Percutaneous Kidney Stone Surgery? One Center Retrospective Study

Burhan Baylan¹,Sercan Sari²,Mehmet Caglar Cakıcı³,Volkan Selmi²,Harun Özdemir⁴,Hakki Ugur Ozok⁵, Ahmet Nihat Karakoyunlu¹,Hikmet Topaloglu¹, Azmi Levent Sagnak¹,Hamit Ersoy¹

¹Department of Urology, University of Health Sciences, Diskapi Yildirim Beyazit Training and Research Hospital,Ankara,Turkey

²Department of Urology, Bozok University Faculty of Medicine,Yozgat,Turkey

³Department of Urology, Medeniyet University, Göztepe Training and Research Hospital,İstanbul,Turkey

⁴Department of Urology,Avcılar State Hospital, İstanbul,Turkey

⁵Department of Urology, Karabük University, Faculty of Medicine,Karabük,Turkey

Keywords: Efficiency, Kidney Stone, Previous Surgery, RIRS, Safety

Abstract

Purpose: In our study, we assessed the efficiency and reliability of retrograde intrarenal surgery secondary to open surgery for kidney stone treatment. Moreover, we compared the efficiency and safety of retrograde intrarenal surgery for the patients with previous history of open surgery, Percutaneous nephrolithotomy, secondary RIRS and primary RIRS.

Materials and Methods: Data was retrospectively reviewed. Patients who had kidney anomalies, who had been stented due to ureteral stricture in the operation and who were < 18 years old, were excluded. There were 30 patients who underwent RIRS secondary to open surgery. The demographic and stone characteristic as well as intraoperative and postoperative data of the patients were recorded. 30 patients with similar demographic and stone characteristics to those patients were selected by match pairing method from patients who had previous PNL, RIRS history and had been undergone primary RIRS . Total 120 patients, in total 4 groups, were included in the study.

Results: Statistically significant difference was detected among the groups with regards to shock wave lithotripsy history and preoperative JJ stent rate, either. There was no statistically significant difference in terms of stone characteristics ,intraoperative and postoperative data

Conclusions: RIRS is an efficient and safe method for kidney stone treatment of the patients with previous history of open surgery, Percutaneous nephrolithotomy and retrograde intrarenal surgery. It has a similar efficiency and safety for the patients who have undergone retrograde intrarenal surgery firstly. This is the first study that compares the patients especially with different previous surgery methods.

Introduction:

Urinary stone disease is a significant health problem affecting human health. Kidney stone prevalence is 1-5% in general⁽¹⁾. Shock Wave Lithotripsy (SWL), percutaneous

nephrolithotomy (PNL), open surgery and recently retrograde intra-renal surgery (RIRS) are used for the treatment of kidney stones. By recently developing technology, certain improvements have been ensured for kidney stone treatment. Non-invasive methods have replaced the invasive ones. Despite significant decrease in preference for open surgery, it is still preferred for selected cases^(2,3). Usage of RIRS has increased thanks to developing technology and increasing experience in recent times. Efficiency of RIRS for kidney stone treatment has been indicated in the studies⁽⁴⁾.

Kidney stone may require repetitive surgical intervention subsequent to surgical treatment. Fibrosis arising after open surgery and changing anatomy may decrease success⁽⁵⁾. There are studies regarding percutaneous nephrolithotomy after open surgery procedures for kidney stone treatment⁽⁶⁻⁸⁾. Although percutaneous nephrolithotomy is an efficient treatment method, life-threatening complications may be observed⁽⁹⁾. There are a limited number of studies regarding usage of RIRS following open surgery^(10,11).

There are studies regarding the factors affecting RIRS success^(12,13). Its usage widened with the advanced technology^(14,15). We want to make the first study especially evaluating the effect of previous stone surgery on RIRS safety and efficacy.

In our study, we assessed the efficiency and reliability of RIRS secondary to open surgery for kidney stone treatment. Moreover, we compared the efficacy of RIRS after previous open surgery, previous PNL and RIRS and primary patients.

Materials and Methods :

The data of the patients who had underwent RIRS in our clinic between 2012-2018 was reviewed retrospectively. To determine the effect of previous surgery history on RIRS safety and efficacy, the records of the patients who had underwent RIRS secondary to open surgery were evaluated. Patients who had kidney anomalies, who had been stented due to ureteral

stricture in the operation and who were < 18 years old, were excluded. There were 30 patients who had undergone RIRS secondary to open surgery. The demographic and stone characteristics as well as intraoperative and postoperative data of the patients were recorded. 30 patients with similar demographic and stone characteristics to those patients were selected by match pairing from patients who had previous PNL, RIRS history and had undergone primary RIRS. The previous open surgery group was divided into groups according to stone size 5-10 mm, 11-15 mm, 16-20 mm, 21-25 mm, 26-30 mm, 31-35 mm, 36-40 mm, 41-45 mm, 46-50mm, 51-55 mm. The same number of procedures were selected randomly from the other groups randomly. The randomization was made similarly for the criteria such as stone laterality, stone number and stone localization. Total 120 patients, in total 4 groups, were included in the study. All patients gave their informed consent for inclusion before they participated in the study. The study was conducted in accordance with the Declaration of Helsinki.

Preoperative routine blood biochemistry tests, complete blood count, coagulation profile and hepatic markers of the patients were analysed. Preoperative urinary cultures of all patients were sterile. Kidney Ureter Bladder Graphy(KUBG), urinary system ultrasonography (US) and unenhanced computerized tomography (CT) were performed preoperatively. The longest stone diameter in imaging was defined as the stone size. In case of multiple stones, total of the longest diameters of each stone was defined as the stone size.

All operations were performed under general anesthesia. Preoperative single dose prophylactic antibiotic treatment was applied for the patients. The ureter was penetrated by placing a hydrophilic guidewire of 0.035/0.038 inch under fluoroscopy guidance by using semi-rigid ureteroscope at modified dorsal lithotomy position. Ureteral access sheath(UAS) (9.5/11.5 F or 11/13 F) (Elite Flex, Ankara, Turkey) was placed into the ureter down to the ureteropelvic junction via guidewire under fluoroscopy guidance. Then, flexible ureteroscope

(Flex-X2, Karl Storz, Tuttlingen, Germany) was moved through the UAS. In case UAS could not be placed, flexible ureteroscope was moved via hydrophilic glidewire and thus access to kidney was ensured. After finding the stone, lithotripsy was applied with 200µm holmium laser (Ho YAG Laser; Dornier MedTech; Munich, Germany / Dornier Med-Tech GmbH, Medilas H20 and HSolvo, Wessling, Germany) thanks to a flexible ureteroscope. The methods of dusting and fragmentation were utilized. All calyces were controlled at the end of the operation. Subsequent to this operation, a JJ stent was placed according to intraoperative conditions. The procedure was completed by placing a 16f foley catheter. In the postoperative first day, the urethral catheter was removed. Following 3 weeks, JJ stent was taken out under daily anaesthesia. Time between starting to endoscopy and JJ stent placing was defined as operation time.

Postoperative control was evaluated by KUBG and US performed on the first postoperative day and unenhanced computerized tomography (CT) performed in the third postoperative month. The patients were followed-up for 3 months. After the controls carried out, patients who were stone free and who had residues <3 mm were accepted as successful. The intraoperative and postoperative data was recorded. The complications were recorded as per Clavien Dindo classification. The groups were compared in terms of efficiency and safety.

Statistical Analysis:

Analyse of data was performed with SPSS for Windows 16.0 package program(SPSS,Chicago).One Sample Kolmogorov Smirnov test was performed to determine whether the distributions of the variables that have numerical values were normal. The distributions of the variables were not normal($p<.05$) except age and body mass index(BMI). The comparisons between the groups were performed with One Way Anova test for the parameters with normal distribution. For the variables with non normal distribution such as stone number, stone size,operation time,scopy time and hospitalisation time, analyse between

the groups were performed with Kruskal Wallis test. The analyse of the nominal variables such as gender, stone laterality, stone localization, UAS usage, postoperative JJ stent, success, SWL history, preoperative JJ stent, anticoagulant usage and complications was performed with pearson chi square test. $P < .05$ value was accepted as statistically significant for the results.

Results:

Total 120 patients were included in our study. Those who had underwent open surgery, PNL, RIRS and primary treatment were divided into Group 1, Group 2, Group 3 and Group 4 respectively.

In terms of demographic data, no statistically significant difference was found out among the groups with regards to the age, sex, BMI and anticoagulant usage. Statistically significant difference was detected among the groups with regards to SWL history and preoperative JJ stent rate, either. ($p < .001$) (Table 1)

In terms of stone data, no statistically significant difference was revealed among the groups with reference to stone laterality, number, size and localization. (Table 1)

In terms of intraoperative data, no statistically significant difference was observed among the groups with regards to average operation and scopy time, use of JJ stent postoperatively and UAS. (Table 2)

In terms of postoperative data, no statistically significant difference was established among the groups with reference to success, hospitalization and complications (Table 2)

Discussion:

Significant changes have occurred in surgical treatment of kidney stone diseases⁽¹⁶⁾. Open surgery has been displaced to the methods such as RIRS and PNL. Stone disease is a morbidity that may show recurrence. It was found out via literature review that previous kidney stone

surgery did not affect success of RIRS⁽¹⁷⁾. There are a limited number of studies regarding success of RIRS after open surgery in the literature. In our study, we aimed to assess the efficiency and safety of RIRS after open surgery that was performed for kidney stone treatment. Moreover, we aimed to compare similar patients who had previous PNL and RIRS histories and those for whom RIRS was applied firstly. This is the first such study in the literature.

In the literature, there are 2 studies which assess RIRS success after open surgery performed for kidney stone treatment. In one of these studies, 53 patients who had underwent RIRS and had an open surgery history for kidney stone were evaluated. In the other study, 38 primary patients with the same characteristics were compared to 32 patients who had underwent RIRS and had an open surgery history for kidney stone. The average operation times were reported to be 79.5 ± 37.8 minutes and 82 minutes^(10,11), respectively. In our study, the average operation time was 47.33 ± 19.33 minutes.

In our study, average stone number and stone size of the patients who had underwent open surgery were 1.93 ± 0.26 and 19.1 ± 11.94 mm, respectively. While average stone number was respectively 3 and 2.7 ± 1.5 in the studies in the literature, average stone size was 14.3mm and 25.4 ± 14.7 mm, respectively.

While the rate of UAS usage was 86.67% in the group that underwent open surgery, the rate of postoperative JJ stent usage was 93.33%. On the other hand, the rate of UAS usage was 77% and 95% and the rate of postoperative JJ stent usage was 100% and 71% respectively in the reviewed studies^(10,11). In terms of success rate, it was 76.67% for the group that had underwent open surgery in our study. In the other studies, it was reported as 79.2% and 82%. Complications were observed in 20.7% and 17% of the patients in the aforementioned studies in the literature^(10,11). In our study, complication was observed in 1 patient in the group that underwent open surgery.

In our study, the patients who had previous open surgery, PNL and RIRS were compared to those who had underwent RIRS firstly. The demographic and stone characteristics of the patients were similar. No statistically significant difference was observed between success and complication rates of the patients. The developments in laser and flexible reoscopes and increasing experience may explain this situation. Similar results were revealed in the studies researching RIRS success and complications, too⁽¹⁸⁻²⁰⁾. Our study is the first one in the literature that compares the patients with similar demographic and kidney stone characteristics who had previous different surgical methods and had underwent RIRS first time.

The limiting factors of our study are its retrospective design and limited number of patients. We need studies designed with larger number of patients and in a prospective design.

Conclusion:

RIRS is an efficient and safe method for kidney stone treatment of the patients with previous history of open surgery, PNL and RIRS. It has a similar efficiency and safety for the patients who had underwent RIRS firstly.

Conflict of Interest:

There is no conflict of interest between the authors.

References

- 1) Menon M, Resnick MI. Urinary Lithiasis: etiology, diagnosis, and medical management. Campbell's Urology, Editor-in-chief: Patrick C. Walsh. Saunders, 2002, 8. Edition, Part 96.
- 2) Honeck P, Wendt NG, Krombach P et al., "Does open stone surgery still play a role in the treatment of urolithiasis? Data of a primary urolithiasis center," *J Endourol*, 2009, 23:1209–1212.

- 3) Paik MI, Resnick MI. Is there a role for open stone surgery? *Urol Clin North Am* 2000, 27(2):323–331
- 4) Wendt NG, Mut T, Krombach P, Michel MS, Knoll T. Do new generation flexible ureterorenoscopes offer a higher treatment success than their predecessors? *Urol Res.* 2011;39:185–8.
- 5) Margel D, Lifshitz DA, Kugel V, Dorfmann D, Lask D, and Livne PM, “Percutaneous nephrolithotomy in patients who previously underwent open nephrolithotomy,” *J Endourol*, 2005,19: 1161–1164.
- 6) Sofikerim M, Demirci D, Gulmez I, Karacagil M, “Does previous open nephrolithotomy affect the outcome of percutaneous nephrolithotomy?” *J Endourol*, 2007,21: 401–403.
- 7) Lojanapiwat B, “Previous open nephrolithotomy: does it affect percutaneous nephrolithotomy techniques and outcome?” *J Endourol*, 2006,20:17–20.
- 8) Gupta R, Gupta A, Singh G, Suri A, Mohan SK, Gupta CL, “PCNL A comparative study in nonoperated and in previously operated (open nephrolithotomy/pyelolithotomy) patients—a single surgeon experience,” *Int Braz J Uro*, 2011,37:739–744.
- 9) Kim SC, Kuo RL, Lingeman JE, “Percutaneous nephrolithotomy: an update,” *Cur Opp in Uro*,2003, 13:235–241.
- 10) Osman, M.M., Gamal, W.M., Gadelmoula, M.M. et al. *Urol Res* ,2012, 40: 403.
<https://doi.org/10.1007/s00240-011-0435-1>
- 11) Alkan E, Saribacak A, Ozkanli AO, Başar MM, Acar O, Balbay MD. Retrograde Intrarenal Surgery in Patients Who Previously Underwent Open Renal Stone Surgery. *Min Inv Sur.* 2015;2015:198765. doi:10.1155/2015/198765.

- 12)** Sari S, Ozok HU, Topaloglu H, et al. The Association of a Number of Anatomical Factors with the Success of Retrograde Intrarenal Surgery in Lower Calyceal Stones. *Urol J.* 2017 Jul 2;14(4):4008-4014.
- 13)** Gucuk A, Yilmaz B, Gucuk S, Uyeturk U. Are Stone Density and Location Useful Parameters That Can Determine the Endourological Surgical Technique for Kidney Stones That Are Smaller than 2 cm? A Prospective Randomized Controlled Trial. *Urol J.* 2018 Aug 26. doi: 10.22037/uj.v0i0.4280. [Epub ahead of print]
- 14)** Javanmard B, Kashi AH, Mazloomfard MM, Ansari Jafari A, Arefanian S . Retrograde Intrarenal Surgery Versus Shock Wave Lithotripsy for Renal Stones Smaller Than 2 cm: A Randomized Clinical Trial. *Urol J.* 2016 Oct 10;13(5):2823-2828.
- 15)** Demirbas A, Yazar VM, Ersoy E, et al. Comparison of Percutaneous Nephrolithotomy and Retrograde Intrarenal Surgery For The Treatment of Multicalyceal and Multiple Renal Stones. *Urol J.* 2018 Nov 17;15(6):318-322. doi: 10.22037/uj.v0i0.4213.
- 16)** Kerbl K, Rehman J, Landman J, Lee D, Sundaram C, Clayman RV. Current management of urolithiasis: progress or regress? *J Endourol* 2002,16:281–288
- 17)** Resorlu B, Unsal A, Gulec H, Oztuna D .A New Scoring System for Predicting Stone-free Rate After Retrograde Intrarenal Surgery: The “Resorlu-Unsal Stone Score” *Urology* ,2012, 80 : 512 - 518
- 18)** Tonyalı S, Yılmaz M, Karaaslan M, Ceylan C, Işıkkay L. Prediction of stone-free status after single-session retrograde intrarenal surgery for renal stones. *Turk J Urol* 2018; 44(6): 473-7.
- 19)** Baş, O., Tuygun, C., Dede, O. et al. *World J Urol* ,2017, 35: 819.
<https://doi.org/10.1007/s00345-016-1930-3>

20) Atis G, Pelit ES, Culpan M, et al. The Fate of Residual Fragments After Retrograde Intrarenal Surgery in Long-Term Follow-up. Urol J. 2018 Jul 23;0(0):4124. doi: 10.22037/uj.v0i0.4124.

Corresponding Author:

Sercan SARI, MD

Bozok University, Department of Urology, Yozgat, Turkey

E-mail: sercansari92@hotmail.com

Phone: +90 5356608838 Fax: + 90 354 2127060

Table 1. Demographic and Stone Characteristics of The Groups

	Group Previous Open Surgery	Group Previous PNL	Group Previous RIRS	Group Primary	<i>p</i>
Age(years)(±SD)	52.16±12.31	49.9±9.84	51.57±11.29	50.5±8.99	.843
Gender(M/F)(n)	19/11	20/10	19/11	18/12	.962
BMI(kg/m ²)(±SD)	29.2±4.6	26.62±3.77	29.76±4.98	27.23±2.54	.056
SWL History (n, %)	9(30)	12(40)	4(13.32)	15(50)	<.001
Anticoagulant Usage (n,%)	0	1(3.33)	0	0	.412
Preoperative JJ Stent (n,%)	2(6.66)	7(23.33)	18(60)	0	<.001
Stone Laterality(R/L)(n)	14/16	15/13	15/13	11/18	.652
Stone Number(n)(±SD)	1.93±0.26	1.96±0.21	1.87±0.15	1.80±0.13	.887
Stone Size(mm) (±SD)	19.1±11.94	19.2±9.15	18.97±5.81	19.67±7.99	.579
Stone Localization (n,%)					.917

Upper Calyx (n,%)	0	0	1(3.33)	1(3.33)	
Lower Calyx (n,%)	12(40)	11(36.67)	12(40)	12(40)	
Mid Calyx (n,%)	3(9.99)	1(3.33)	3(9.99)	3(9.99)	
Pelvis (n,%)	7(23.33)	4(13.32)	7(23.33)	7(23.33)	
Multicaliceal (n,%)	6(20)	10(33.33)	5(16.65)	5(16.65)	
Proximal Ureter(n,%)	2(6.66)	4(13.32)	2(6.66)	2(6.66)	

SD:Standart Deviation, **M:**Male, **F:**Female, **BMI:**Body Mass Index, **SWL:**Shock Wave Lithotripsy

Mm:Milimeter, **R:**Right, **L:** Left , **PNL:** Percutaneous Nephrolithotomy, **RIRS:** Retrograde Intrarenal Surgery

Table 2 Intraoperative and Postoperative Data of The Groups

	Group Previous Open Surgery	Group Previous PNL	Group Previous RIRS	Group Primary	<i>p</i>
Average Operation Time(min.) (±SD)	47.33±19.33	56.67±28.81	61.67±27	52.53±14.84	.157
Average Scopy Time (Sc.) (±SD)	26.17±18.08	56.40±18.25	32.2±26.73	39.4±29.19	.170
Postoperative JJ stent, n (%)	28(93.33)	28(93.33)	30(100)	25(83.33)	.106
Ureteral Access Sheath Usage , n (%)	26(86.67)	24(80)	27(90)	29(96.67)	.240
Average Hospitalisation Time(±SD) (day)	1	1	2,55±1.47	1	.392
Success , (n) (%)	23(76.67)	19(63.33)	19(63.33)	24(80)	.341
Stone-free , (n) (%)	23(76.67)	18(60)	18(60)	21(70)	
Residuel fragment (<3mm) , (n) (%)	0	1(3.33)	1(3.33)	3(10)	

Residuel fragment (≥3mm), (n) (%)	7(23.33)	11(36.67)	11(36.67)	6(20)	
Complication rate , n (%)	1(3.33)	7(23.33)	3(10)	4(13.32)	.126
Intraoperative Complication, (n) (%)	1(3.33)	2(6.66)	1(3.33)	3(10)	
Mucosal Injury, n (%)	1(3.33)	1(3.33)	1(3.33)	3(10)	
Bleeding , n (%)	0	1(3.33)	0	0	
Postoperative Complication , n (%)	0	7(23.33)	2(6.66)	1(3.33)	
Fever (Clavien I) , n (%)	0	7(23.33)	0	0	
Bleeding (Clavien I) , n (%)	0	4(13.32)	0	0	
Urinary Tract Infection (Clavien II) , n (%)	0		0	1(3.33)	
Perirenal Hematom(Clavien 3a), n(%)	0	1(3.33)	0	0	
Steinstrasse(Clavien IIIb), n (%)	0	2(6.66)	2(6.66)	0	

Min:Minute, **Sc**: Second, **SD**:Standart Deviation, **MM**:Milimeter, , **PNL**: Percutaneous Nephrolithotomy, **RIRS**: Retrograde Intrarenal Surger